arXiv:1112.2122v1 [math.AP] 9 Dec 2011

A TWO DIMENSIONAL ADLER-MANIN TRACE AND BI-SINGULAR OPERATORS

FARZAD FATHIZADEH, MASOUD KHALKHALI, FABIO NICOLA, LUIGI RODINO

Abstract. Motivated by the theory of bi-singular pseudodifferential operators, we find an algebraic setting for a noncommutative residue defined first analytically for such operators.

1. Introduction

The Adler-Manin trace was discovered for one dimensional pseudodifferential symbols [1, 13]. It was generalized by Wodzicki to the algebra of pseudodifferential operators on a manifold of arbitrary dimension [22]. A vast generalization of Wodzicki's noncommutative residue was obtained by Connes and Moscovici for spectral triples in the context of the local index formula in noncommutative geometry [5]. Also, a twisted version of the Adler-Manin trace was studied in [9].

In this paper, motivated by the theory of bi-singular pseudodifferential operators [17], we find an algebraic setting for a noncommutative residue defined analytically in [14]. Fredholm properties, index and residue were studied for an algebra of pseudodifferential operators, denoted by $HL(X_1 \times X_2)$, on the product of two manifolds X_1 and X_2 [14]. Considering the case $X_1 = X_2 = \mathbb{S}^1$, and writing I for the two sided ideal of the operators which are smoothing with respect to at least one of the variables on $\mathbb{T} = \mathbb{S}^1 \times \mathbb{S}^1$, we get an isomorphism

$$HL(\mathbb{T})/I \to \Psi(A, \delta_1, \delta_2), \tag{1}$$

where A is the direct sum of 4 copies of the algebra $C^\infty(\mathbb{T})$ and δ_1, δ_2 are the standard partial derivatives. Each element in the right-hand side of (1) has then a representation of the form

$$\sum_{m=-\infty}^{M} \sum_{n=-\infty}^{N} a_{m,n} \xi_1^m \xi_2^n, \tag{2}$$

with $a_{m,n} = (a_{m,n}^{(s,t)})_{s,t=1,2}$. We have for $a \in A$ the linearly independent traces

$$\tau_{s,t}(a) = \iint_{\mathbb{T}} a^{(s,t)}(x_1, x_2)\, dx_1\, dx_2, \qquad s, t = 1, 2,$$

and correspondingly 4 residues are defined by $\tau_{s,t}(a_{-1,-1})$.

Bi-singular operators P were originally studied, during the years 60's and 70's, under the form

$$Pf(z_1, z_2) = b_0(z_1, z_2) f(z_1, z_2) + \frac{1}{\pi i} \int_{\mathbb{S}^1} \frac{b_1(z_1, z_2, \zeta_1)}{\zeta_1 - z_1} f(\zeta_1, z_2)\, d\zeta_1 \tag{3}$$

$$+ \frac{1}{\pi i} \int_{\mathbb{S}^1} \frac{b_2(z_1, z_2, \zeta_2)}{\zeta_2 - z_2} f(z_1, \zeta_2)\, d\zeta_2 - \frac{1}{\pi^2} \iint_{\mathbb{S}^1 \times \mathbb{S}^1} \frac{b_{1,2}(z_1, z_2, \zeta_1, \zeta_2)}{(z_1 - \zeta_1)(z_2 - \zeta_2)} f(\zeta_1, \zeta_2) d\zeta_1\, d\zeta_2,$$

where we regard $\mathbb{S}^1$ as the unit circle in the complex plane and we understood counter-clockwise integration in the Cauchy sense. We assume the functions $b_0, b_1, b_2, b_{1,2}$ are C^∞ with respect to all of the variables. These operators, bounded on $L^2(\mathbb{T})$, can be regarded as operators of order zero in $HL(\mathbb{T})$, namely $a_{m,n} = 0$ for $(m, n) \neq (0, 0)$ in (2). To respect the algebraic isomorphism in (1), one sets in (2) $a_{0,0} = (a_{0,0}^{(s,t)})_{s,t=1,2}$ with

$$\begin{aligned} a_{0,0}^{(s,t)} = b_0(z_1, z_2) - (-1)^s b_1(z_1, z_2, z_1) - (-1)^t b_2(z_1, z_2, z_2) \\ + (-1)^{s+t} b_{1,2}(z_1, z_2, z_1, z_2), \end{aligned}$$

with $z_j = e^{2\pi i x_j}$, $j = 1, 2$.

The literature in this connection is wide, involving problems of operator theory, harmonic analysis and several complex variables, see for example [23, 21, 10], coming from the school of A. Zygmund, and [19, 15, 16, 11], from the school of B. Simonenko.

Starting from the 70's, bi-singular operators were reconsidered from the point of view of the theory of pseudodifferential operators, see [8] and [17], emphasizing the connection with the proof of the index theorem [3]. Roughly: the algebra $HL(\mathbb{T})$ is obtained by composing P in (3) with partial derivatives on $\mathbb{T}$ and their inverses. This originates terms of positive and negative orders in (2), and gives rise to non-trivial residues.

A natural question is whether a related class of pseudodifferential operators exists, providing in (1) an isomorphism with $\Psi(A, \delta_1, \delta_2)$ where simply $A = C^\infty(\mathbb{T})$. To this end, we may address to the Toeplitz operators in a quadrant of [6, 7, 4, 20].

This paper is organized as follows. In Section 2 we recall some standard notions and constructions, and the Adler-Manin trace in the one dimensional case. In Section 3, starting from an algebra A equipped with a trace which is invariant with respect to two derivations of A, we construct a two dimensional algebra of formal pseudodifferential symbols. Then, we define a noncommutative residue on this algebra and prove that it is a trace functional. In Section 4, as an example of our two dimensional analogue of the Adler-Manin noncommutative residue, we shall consider the residue of the bi-singular operators, already treated in [14].

2. The Adler-Manin Trace

Let A be a unital associative algebra over $\mathbb{C}$ which is not necessarily commutative. Recall that a *derivation* on A is a linear map $\delta : A \to A$ such that

$$\delta(ab) = a\delta(b) + \delta(a)b$$

for all $a, b \in A$. Given a pair (A, δ) as above, the algebra of *formal differential symbols* $D(A, \delta)$ [12, 13] is, by definition, the algebra generated by A and a symbol ξ subject to the relations

$$\xi a - a\xi = \delta(a) \tag{4}$$

for all $a \in A$. There is a unique expression of the form

$$D = \sum_{i=0}^{N} a_i \xi^i, \qquad N \geq 0, \qquad a_i \in A,$$

for every element of $D(A,\delta)$. We think of D as a differential operator of order at most N. Using (4), one can prove by induction that

$$\xi^n a = \sum_{j=0}^{n} \binom{n}{j} \delta^j(a)\xi^{n-j} \tag{5}$$

for all $a \in A$ and $n \geq 0$. Using (5), we obtain the following multiplication formula in $D(A,\delta)$:

$$\Big(\sum_{i=0}^{M} a_i\xi^i\Big)\Big(\sum_{j=0}^{N} b_j\xi^j\Big) = \sum_{n=0}^{M+N}\Big(\sum_{i,j,k}\binom{i}{k} a_i\delta^k(b_j)\Big)\xi^n$$

where the internal summation is over all $0 \leq k \leq i \leq M$, and $0 \leq j \leq N$ such that $i+j-k=n$.

By formally inverting ξ in $D(A,\delta)$ and completing the resulting algebra, one obtains the algebra of *formal pseudodifferential symbols* of (A,δ) [1, 13, 12], denoted by $\Psi(A,\delta)$. More precisely, it is defined as follows. Elements of $\Psi(A,\delta)$ consist of formal sums

$$D = \sum_{i=-\infty}^{N} a_i\xi^i$$

with $a_i \in A$, and $N \in \mathbb{Z}$. Its multiplication is defined by first extending (5) to all $n \in \mathbb{Z}$:

$$\xi^n a = \sum_{j=0}^{\infty}\binom{n}{j}\delta^j(a)\xi^{n-j}.$$

Here the binomial coefficient $\binom{n}{j}$ for $n \in \mathbb{Z}$, and non-negative $j \in \mathbb{Z}$, is defined by $\binom{n}{j} := \frac{n(n-1)\cdots(n-j+1)}{j!}$. Notice that for $n<0$, we have an infinite formal sum. In general, for $D_1 = \sum_{i=-\infty}^{M} a_i\xi^i$, and $D_2 = \sum_{j=-\infty}^{N} b_j\xi^j$, the multiplication is defined by

$$D_1D_2 = \sum_{n=-\infty}^{M+N}\Big(\sum_{i,j,k}\binom{i}{k}a_i\delta^k(b_j)\Big)\xi^n$$

where the internal summation is over all integers $i \leq M$, $j \leq N$, and $k \geq 0$ such that $i+j-k=n$.

Now consider a δ-*invariant trace* $\tau : A \to \mathbb{C}$. By definition, τ is a linear map and we have

$$\tau(ab) = \tau(ba), \qquad \tau(\delta(a)) = 0,$$

for all $a, b \in A$. The *Adler-Manin noncommutative residue* [1, 13, 12], res : $\Psi(A,\delta) \to \mathbb{C}$ is defined by

$$\operatorname{res}\Big(\sum_{i=-\infty}^{N} a_i\xi^i\Big) = \tau(a_{-1}).$$

One checks that res is indeed a trace, i.e.

$$\operatorname{res}[D_1, D_2] = 0$$

for all $D_1, D_2 \in \Psi(A,\delta)$. Equivalently, one shows that the map res : $\Psi(A,\delta) \to A/([A,A] + \operatorname{im}\delta)$

$$D \mapsto a_{-1} \qquad \operatorname{mod} \qquad [A,A] + \operatorname{im}\delta$$

is a trace on $\Psi(A,\delta)$ with values in $A/([A,A]+\operatorname{im}\delta)$.

A relevant example is when $A=C^\infty(\mathbb{S}^1)$, the algebra of smooth functions on the circle $\mathbb{S}^1=\mathbb{R}/\mathbb{Z}$, with

$$\tau(f)=\int_0^1 f(x)\,dx, \qquad \delta(f)=f'$$

for all $f\in C^\infty(\mathbb{S}^1)$. Then the noncommutative residue coincides with Wodzicki residue on the algebra of classical pseudodifferential operators on the circle, see Section 4 below.

3. Two Dimensional Noncommutative Residue

In this section we extend the Adler-Manin trace to a two dimensional case where an algebra A equipped with two derivations $\delta_1,\delta_2:A\to A$ is considered. Assuming that $\delta_1\delta_2=\delta_2\delta_1$, we define an algebra of formal pseudodifferential symbols, denoted by $\Psi(A,\delta_1,\delta_2)$ whose elements are formal series of the form

$$\sum_{m=-\infty}^{M}\sum_{n=-\infty}^{N} a_{m,n}\xi_1^m\xi_2^n, \qquad a_{ij}\in A, \qquad M,N\in\mathbb{Z}.$$

The multiplication of this algebra is essentially defined by the following relations:

$$\xi_1\xi_2=\xi_2\xi_1,$$
$$\xi_i\xi_i^{-1}=\xi_i^{-1}\xi_i=1,$$
$$\xi_i a=a\xi_i+\delta_i(a), \qquad a\in A,$$

for $i=1,2$. Similar to the one dimensional case described in Section 2, for any integer n, we define

$$\xi_i^n a=\sum_{j=0}^{\infty}\binom{n}{j}\delta^j(a)\xi_i^{n-j}, \qquad a\in A, \qquad i=1,2. \tag{6}$$

Using these relations, we can write the following explicit formula for the multiplication in $\Psi(A,\delta_1,\delta_2)$:

$$\Big(\sum_{m=-\infty}^{M}\sum_{n=-\infty}^{N} a_{m,n}\xi_1^m\xi_2^n\Big)\Big(\sum_{m'=-\infty}^{M'}\sum_{n'=-\infty}^{N'} b_{m',n'}\xi_1^{m'}\xi_2^{n'}\Big)$$
$$=\sum_{m=-\infty}^{M}\sum_{n=-\infty}^{N}\sum_{m'=-\infty}^{M'}\sum_{n'=-\infty}^{N'}\sum_{i,j=0}^{\infty}\binom{m}{i}\binom{n}{j}a_{m,n}\delta_1^i\delta_2^j(b_{m',n'})\xi_1^{m+m'-i}\xi_2^{n+n'-j}.$$

In the following, starting from an invariant trace $\tau:A\to\mathbb{C}$, we define a noncommutative residue on $\Psi(A,\delta_1,\delta_2)$ and prove that it is a trace functional on this algebra. First we prove a lemma:

Lemma 3.1. *Let $\delta:A\to A$ be a derivation and $\tau:A\to\mathbb{C}$ be a linear functional such that $\tau\circ\delta=0$. Then for any $a,b\in A$ and non-negative integer i, we have*

$$\tau(\delta^i(a)b)=(-1)^i\tau(a\delta^i(b)).$$

Proof. For any $a, b \in A$, we have

$$\tau(\delta(ab)) = \tau(\delta(a)b) + \tau(a\delta(b)) = 0.$$

Therefore

$$\tau(\delta(a)b) = -\tau(a\delta(b)),$$

from which the lemma follows easily. □

Now we define a noncommutative residue and prove the main theorem:

Theorem 3.2. *Let* $\tau : A \to \mathbb{C}$ *be a trace such that* $\tau \circ \delta_i = 0$ *for* $i = 1, 2$. *Then the linear functional* $\mathrm{Res} : \Psi(A, \delta_1, \delta_2) \to \mathbb{C}$ *defined by*

$$\mathrm{Res}(\sum_{m=-\infty}^{M} \sum_{n=-\infty}^{N} a_{m,n}\xi_1^m \xi_2^n) = \tau(a_{-1,-1})$$

is a trace on $\Psi(A, \delta_1, \delta_2)$.

Proof. Let $a, b \in A$ and $m, n, m', n' \in \mathbb{Z}$. It suffices to show that

$$\mathrm{Res}(a\xi_1^m \xi_2^n b\xi_1^{m'} \xi_2^{n'}) = \mathrm{Res}(b\xi_1^{m'} \xi_2^{n'} a\xi_1^m \xi_2^n).$$

On the left side of the latter we have:

$$\begin{aligned}
a\xi_1^m \xi_2^n b\xi_1^{m'} \xi_2^{n'} &= a\xi_1^m \sum_{i=0}^{\infty} \binom{n}{i} \delta_2^i(b)\xi_2^{n-i}\xi_1^{m'}\xi_2^{n'} \\
&= a\sum_{i=0}^{\infty} \binom{n}{i} \xi_1^m \delta_2^i(b)\xi_1^{m'}\xi_2^{n'+n-i} \\
&= a\sum_{i=0}^{\infty} \binom{n}{i} \sum_{j=0}^{\infty} \binom{m}{j} \delta_1^j\delta_2^i(b)\xi_1^{m-j}\xi_1^{m'}\xi_2^{n+n'-i} \\
&= \sum_{i=0}^{\infty}\sum_{j=0}^{\infty} \binom{n}{i}\binom{m}{j} a\delta_1^j\delta_2^i(b)\xi_1^{m+m'-j}\xi_2^{n+n'-i}.
\end{aligned}$$

Therefore

$$\mathrm{Res}(a\xi_1^m \xi_2^n b\xi_1^{m'} \xi_2^{n'}) = \tau\Big(\sum \binom{n}{i}\binom{m}{j} a\delta_1^j\delta_2^i(b)\Big), \tag{7}$$

where the summation is over all non-negative integers i, j such that $m+m'-j = -1$ and $n + n' - i = -1$.

Similarly we have:

$$b\xi_1^{m'} \xi_2^{n'} a\xi_1^m \xi_2^n = \sum_{i=0}^{\infty}\sum_{j=0}^{\infty} \binom{m'}{j}\binom{n'}{i} b\delta_1^j\delta_2^i(a)\xi_1^{m+m'-1}\xi_2^{n+n'-1}.$$

Therefore

$$\begin{aligned}
\mathrm{Res}(b\xi_1^{m'} \xi_2^{n'} a\xi_1^m \xi_2^n) &= \tau\Big(\sum \binom{m'}{j}\binom{n'}{i} b\delta_1^j\delta_2^i(a)\Big) \\
&= \tau\Big(\sum \binom{m'}{j}\binom{n'}{i} \delta_1^j\delta_2^i(a)b\Big) \\
&= (-1)^{j+i}\tau\Big(\sum \binom{m'}{j}\binom{n'}{i} a\delta_2^i\delta_1^j(b)\Big),
\end{aligned}$$

where again the summation is over all non-negative integers i, j such that $m+m'-j=-1$ and $n+n'-i=-1$. Note that in the last two equalities, we have used the trace property of τ and Lemma 3.1 respectively. Therefore, in order to show that (7) and (8) are equal, it suffices to show that if i, j are fixed non-negative integers such that $m+m'-j=-1$ and $n+n'-i=-1$, then

$$\binom{m}{j}\binom{n}{i} = (-1)^{j+i}\binom{m'}{j}\binom{n'}{i}.$$

This is easy to see since

$$\begin{aligned}
(-1)^i\binom{n'}{i} &= (-1)^i\binom{-n+i-1}{i} \\
&= (-1)^i\frac{(-n+i-1)(-n+i-1-1)\cdots(-n+i-1-i+1)}{i!} \\
&= \frac{(n-i+1)(n-i+2)\cdots(n)}{i!} \\
&= \binom{n}{i},
\end{aligned}$$

and similarly

$$(-1)^j\binom{m'}{j} = \binom{m}{j}.$$

□

Example 3.1. Let $A = C^\infty(\mathbb{T})$, $\delta_1 = \frac{\partial}{\partial\theta_1}$, $\delta_2 = \frac{\partial}{\partial\theta_2}$, and

$$\tau(f) = \int_0^1\int_0^1 f(\theta_1,\theta_2)\,d\theta_1\,d\theta_2.$$

Then

$$\operatorname{Res}\Big(\sum_{m=-\infty}^{M}\sum_{n=-\infty}^{N} a_{m,n}\xi_1^m\xi_2^n\Big) = \int_0^1\int_0^1 a_{-1,-1}(\theta_1,\theta_2)\,d\theta_1\,d\theta_2.$$

4. SINGULAR AND BI-SINGULAR OPERATORS

We add here some detail concerning the bi-singular operators mentioned in the introduction. To be definite, we consider first the one dimensional case.

Singular integral operators [2, 18] (Classical pseudodifferential operators of order $N \in \mathbb{Z}$ on $\mathbb{S}^1 = \mathbb{R}/\mathbb{Z}$). Classical pseudodifferential operators on the circle can be defined by a global symbol $a(x,\xi)$, $x \in \mathbb{R}/\mathbb{Z}$, $\xi \in \mathbb{R}$, with an asymptotic expansion in homogeneous terms. Modulo regularizing operators, i.e. mapping $\mathcal{D}'(\mathbb{S}^1) \to C^\infty(\mathbb{S}^1)$, this algebra can be therefore identified with that of formal series

$$\sigma = \sum_{n=-\infty}^{N} a_{n,1}(x)\xi_+^n + a_{n,2}(x)\xi_-^n, \tag{8}$$

where $\xi_+ = \max\{\xi, 0\}$, $\xi_- = \min\{\xi, 0\}$.

This corresponds to the Adler-Manin algebra $\Psi(A,\delta)=\Big\{\sum_{n=-\infty}^{N}a_n\xi^n\Big\}$, with $A=C^\infty(\mathbb{S}^1)\bigoplus C^\infty(\mathbb{S}^1)$, $\delta=\frac{d}{dx}$, $a_n=(a_{n,1},a_{n,2})$. There are two linearly independent traces on A, namely

$$\tau_s(a)=\int_0^1 a^{(s)}(x)\,dx,\qquad s=1,2,\qquad a=(a^{(1)},a^{(2)})\in A,$$

and therefore two residues

$$\operatorname{Res}_s\Big(\sum_{n=-\infty}^{N}a_n\xi^n\Big)=\tau_s(a_{-1})=\int_0^1 a_{n,s}(x)\,dx,\qquad s=1,2$$

which are known as Wodzicki's residues.

Introducing on $\Psi(A,\delta)$ the Toeplitz projection

$$P_+\sigma=\sum_{n=-\infty}^{N}a_{n,1}(x)\xi_+^n,$$

i.e. setting $a_{n,2}=0$ for all n in (8), we may define the quotient subalgebra of the asymptotic expansions of the classical Toeplitz operators, cf. the example at the end of Section 2, where A is simply given by $C^\infty(\mathbb{S}^1)$.

Bi-singular integral operators. We consider the so-called algebra of bi-singular integral operators on $\mathbb{T}=\mathbb{S}^1\times\mathbb{S}^1$, which are defined by a global symbol

$$a(x_1,x_2;\xi_1,\xi_2),$$

where $x_1,x_2\in\mathbb{R}/\mathbb{Z}$, $\xi_1,\xi_2\in\mathbb{R}$, with an asymptotic expansion in terms positively homogeneous with respect to ξ_1 and ξ_2 separately. Modulo sums of operators regularizing in x_1 and, respectively, x_2, they are identified with formal series

$$\sigma=\sum_{m=-\infty}^{M}\sum_{n=-\infty}^{N}\sigma_{m,n},\qquad\text{with}\qquad\sigma_{m,n}=\sum_{s,t=1,2}a_{m,n}^{(s,t)}(x_1,x_2)\xi_{1,s}^m\xi_{2,t}^n,$$

where $\xi_{j,1}=\max\{\xi_j,0\}$, $\xi_{j,2}=\min\{\xi_j,0\}$, $j=1,2$.

This corresponds to the algebra

$$\Psi(A,\delta_1,\delta_2)=\Big\{\sum_{m=-\infty}^{M}\sum_{n=-\infty}^{N}a_{m,n}\xi_1^m\xi_2^n\Big\}$$

considered in Section 3, with

$$A=C^\infty(\mathbb{T})^4,\qquad\delta_1=\frac{\partial}{\partial x_1},\qquad\delta_2=\frac{\partial}{\partial x_2},\qquad a_{m,n}=(a_{m,n}^{(s,t)})_{s,t=1,2}\in A.$$

Now we have four linearly independent traces on A, namely

$$\tau_{s,t}(a)=\int_0^1\int_0^1 a^{(s,t)}(x_1,x_2)\,dx_1\,dx_2,\qquad a=(a^{(s,t)})\in A,\qquad s,t=1,2,$$

and therefore four residues

$$\operatorname{Res}_{s,t}\Big(\sum_{m=-\infty}^{M}\sum_{n=-\infty}^{N}a_{m,n}\xi_1^j\xi_2^k\Big)=\tau_{s,t}(a_{-1,-1}),\qquad s,t=1,2.$$

They correspond to those found in [14] for general bi-singular operators on the product of two manifolds.

Note finally that the projection

$$P_{++}\sigma = \sum_{m=-\infty}^{M} \sum_{n=-\infty}^{N} a_{m,n}^{(1,1)}(x_1, x_2)\xi_{1,1}^{m}\xi_{2,1}^{n}$$

gives the algebra of the asymptotic expansions for Toeplitz operators in the quarter plane, corresponding to Example 3.1.

References

[1] M. Adler, On a trace functional for formal pseudo differential operators and the symplectic structure of the Korteweg-de Vries type equations. Invent. Math. 50 (1978/79), no. 3, 219–248.

[2] M.S. Agranovich, Spectral properties of elliptic pseudodifferential operators on a closed curve. (Russian) Funktsional Anal. i Priložen 13 (1979), no. 4, 54–56. (English translation in Functional Analysis and Its applications. 13, p. 279–281).

[3] M. F. Atiyah and I. M. Singer, The index of elliptic operators. Ann. of Math. 87 (1968), 484–530.

[4] L. A. Coburn, R. G. Douglas and I. Singer, An index theorem for Wiener-Hopf operators on the discrete quarter-plane. J. Differential Geometry 6 (1972), 587-593.

[5] A. Connes and H. Moscovici, The local index formula in noncommutative geometry. Geom. Funct. Anal. 5 (1995), no. 2, 174–243.

[6] R. G. Douglas and R. Howe, On the C^*-algebras of Toeplitz operators on the quarter plane. Trans. Amer. Math. Soc. 158 (1971), 203–217.

[7] R. G. Douglas, On the invertibility of a class of Toeplitz operators on the quarter-plane. Indiana Univ. Math. J. 21 (1971/1972), 1031–1035.

[8] R. V. Dudučava, On bisingular integral operators with discontinuous coefficients. Mat. Sbornik 101 (1976); transl. Math. USSR Sbornik 30 (1976), 515–537.

[9] F. Fathizadeh and M. Khalkhali, The algebra of formal twisted pseudodifferential symbols and a noncommutative residue. Letters in Mathematical Physics (2010), 94:41–61.

[10] C. Fefferman, Estimates for double Hilbert transforms. Studia Math. 44 (1972), 1–15.

[11] V.A. Kakičev. Regularization and pseudoregularization of complete bisingular operators with Cauchy kernels and normal characteristic part. Dokl. Akad. Nauk SSSR 229 (1976), 19–22; transl. Soviet Math Dokl. 17 (1976), 952–956.

[12] Ch. Kassel, Le résidu non commutatif (d'après M. Wodzicki). (French) [The noncommutative residue (after M. Wodzicki)] Séminaire Bourbaki, Vol. 1988/89. Astérisque No. 177-178 (1989), Exp. No. 708, 199–229.

[13] Ju. I. Manin, Algebraic aspects of nonlinear differential equations. (Russian) Current problems in mathematics, Vol. 11 (Russian), pp. 5–152. (errata insert) Akad. Nauk SSSR Vsesojuz. Inst. Naučn. i Tehn. Informacii, Moscow, 1978.

[14] F. Nicola and L. Rodino, Residue index for bisingular operators. In "C^* algebras and Elliptic Theory", Trends in Mathematics (2006), 187–202.

[15] V. S. Pilidi, The index of bisingular operators. Funkcional Anal. i Priložen 7 (1973), 93–94; transl. Functional Anal. Appl. 7 (1973), 337–338.

[16] V. S. Pilidi, L. I. Sazonov, A priori estimates for characteristic bisingular integral operators. Dokl. Akad. Nauk SSSR 217 (1974), 285–287; transl. Soviet Math. Dokl. 15 (1974), 1064–1067.

[17] L. Rodino, A class of pseudodifferential operators on the product of two manifolds and applications. Ann. Scuola Norm. Sup. Pisa 2 ser. IV (1975), 287–302.

[18] M. Ruzhansky and V. Turunen, pseudodifferential operators and symmetries. Pseudodifferential Operators, Theory and Applications, Birkhäuser, Basel, 2010.

[19] I. B. Simonenko, On the question of solvability of bisingular and polysingular equations. Funkcional Anal. i Priložen 5 (1971), 93–94; transl. Functional Anal. Appl. 5 (1971), 81–83.

[20] G. Strang, Toeplitz operators in a quarter-plane. Bull. Amer. Math. Soc. 76 (1970), 1303–1307.

[21] T. Walsh, On the existence of double singular integrals for kernels without smoothness. Proc. Amer. Math. Soc. 28 (1971), 439–445.

[22] M. Wodzicki, Noncommutative residue. I. Fundamentals. K-theory, arithmetic and geometry (Moscow, 1984–1986), 320–399, Lecture Notes in Math., 1289, Springer, Berlin, 1987.

[23] A. Zygmund, On the boundary values of functions of several complex variables I. Fund. Math. 36 (1949), 207–235.

F. Fathizadeh: Department of Mathematics and Statistics, York University, Toronto, Ontario, Canada, M3J 1P3

M. Khalkhali: Department of Mathematics, The University of Western Ontario, London, Ontario, Canada, N6A 5B7

F. Nicola: Dipartimento di Matematica, Politecnico di Torino, Corso Duca degli Abruzzi 24, 10129 Torino, Italy

L. Rodino: Dipartimento di Matematica, Università degli Studi di Torino, Via Carlo Alberto 10, 10123 Torino, Italy